

OUTER COMMUTATOR WORDS ARE UNIFORMLY CONCISE

GUSTAVO A. FERNÁNDEZ-ALCOBER AND MARTA MORIGI

ABSTRACT. We prove that outer commutator words are uniformly concise, i.e. if an outer commutator word ω takes m different values in a group G , then the order of the verbal subgroup $\omega(G)$ is bounded by a function depending only on m , and not on ω or G . This is obtained as a consequence of a structure theorem for the subgroup $\omega(G)$, which is valid if G is soluble, and without assuming that ω takes finitely many values in G . More precisely, there is an abelian series of $\omega(G)$, such that every section of the series can be generated by values of ω all of whose powers are also values of ω in that section. For the proof of this latter result, we introduce a new representation of outer commutator words by means of binary trees, and we use the structure of the trees to set up an appropriate induction.

1. INTRODUCTION

Let X be a set of symbols, to which we refer as *indeterminates*. In group theory, a word ω over X is an element of the free group having X as a free basis. If the expression for ω involves k different indeterminates, then for every group G , we obtain a function from G^k to G by substituting group elements for the indeterminates. Thus we can consider the *set* G_ω of all values taken by this function, that is,

$$G_\omega = \{\omega(g_1, \dots, g_k) \mid g_i \in G \text{ for all } i = 1, \dots, k\}.$$

The subgroup generated by G_ω is called the *verbal subgroup* of ω in G , and is denoted by $\omega(G)$. We say that two words are *equivalent* if they can be transformed into each other by simply changing the names of the indeterminates. Obviously, equivalent words define the same set of values and the same verbal subgroup. For this reason, we may assume if necessary that all words are defined over the countable set $X = \{x_1, x_2, \dots\}$, and that if ω involves k indeterminates, these are given by the symbols $x_1, \dots, x_k$.

Words which are formed by taking commutators are particularly interesting. Among them, we have the *lower central words* γ_i , on i indeterminates, which are given by

$$\gamma_1 = x_1, \quad \gamma_i = [\gamma_{i-1}, x_i] = [x_1, \dots, x_i], \quad \text{for } i \geq 2,$$

2000 *Mathematics Subject Classification.* Primary 20F10; Secondary 12L10.

Key words and phrases. Outer commutator words; Verbal subgroups; Ultraproducts.

The first author is supported by the Spanish Ministry of Science and Innovation, grant MTM2008-06680-C02-02, partly with FEDER funds, and by the Basque Government, grant IT-252-07. The second author is partially supported by MIUR (Project “Teoria dei Gruppi e applicazioni”).

and the *derived words* δ_i , on 2^i indeterminates, defined recursively by

$$\delta_0 = x_1, \quad \delta_i = [\delta_{i-1}(x_1, \dots, x_{2^{i-1}}), \delta_{i-1}(x_{2^{i-1}+1}, \dots, x_{2^i})], \quad \text{for } i \geq 1.$$

The words γ_i and δ_i are particular instances of *outer commutator words*, which are words obtained by nesting commutators, but using always *different indeterminates*. Thus $[[x_1, x_2], [x_3, x_4, x_5], x_6]$ is an outer commutator word, but the Engel word $[x_1, x_2, x_2, x_2]$ is not.

A word ω is said to be *concise* if, for every group G , the finiteness of the set G_ω implies that of $\omega(G)$. In the 1960's, Turner-Smith published a couple of papers [7, 8] related to word values and verbal subgroups, where he indicates that Philip Hall had conjectured that every word is concise, and that Hall himself had proved this for every non-commutator word (i.e. a word outside the commutator subgroup of the free group), and for lower central words. In [8], Turner-Smith showed that also derived words are concise, and Jeremy Wilson [9] subsequently extended this result to all outer commutator words. On the other hand, Hall's conjecture was eventually refuted in 1989 by Ivanov, see [3].

If a word ω is concise, it is natural to ask whether conciseness can be expressed in a quantitative form; more precisely, provided that $|G_\omega| = m$, can we bound $|\omega(G)|$ by a function depending only on m ? The answer to this question is positive, but this does not seem to be widely known among group theorists and, to the best of our knowledge, there is no reference in the literature containing this result. For this reason, we have included an appendix at the end of the paper in which we give two different proofs of this fact. Both proofs need the ultraproduct construction for groups, over a non-principal ultrafilter. The first one uses Łoś's Theorem from model theory, while the second one is derived directly from the definition of an ultraproduct, and is due to Avinoam Mann. Note that the existence of non-principal ultrafilters is independent of the Zermelo-Fraenkel (ZF) axioms for set theory; it can be proved by using the Axiom of Choice (but is not equivalent to it).

The ultraproduct argument only shows the existence of bounds for concise words, but it does not provide any explicit expressions for these bounds. In the case of the commutator $\gamma_2 = [x_1, x_2]$, one can use the results bounding the order of the derived subgroup G' in terms of the breadth (maximum size of a conjugacy class) of G . If G contains at most m commutators, it follows that:

- (i) If G is soluble, then $|G'| \leq m^{\frac{1}{2}(5+\log_2 m)}$. (P. Neumann and Vaughan-Lee, [4].)
- (ii) For a general group, $|G'| \leq m^{\frac{1}{2}(13+\log_2 m)}$. (Segal and Shalev, [6].)

More recently, Brazil, Krasilnikov and Shumyatsky [1] have given explicit bounds for all lower central words and for all derived words; as a matter of fact, they find a single upper bound for this infinite family of words, namely $(m!)^m$. The first main result of this paper shows that an even better uniform bound applies to all outer commutator words.

Theorem A. *Let ω be an outer commutator word and let G be a group. If $|G_\omega| = m$, then:*

- (i) *If G is soluble, $|\omega(G)| \leq 2^{m-1}$.*

- (ii) If G is not soluble, $|\omega(G)| \leq (m-1)^{m-1}$.

We suspect that the bounds of Theorem A might be sharpened to get close to the bounds given above for the word γ_2 . Nevertheless, an examination of the papers giving upper bounds for $|G'|$ in terms of the breadth clearly suggests that this would be better the subject of an independent paper, devoted specifically to this question. For this reason, we have not attempted to obtain sharp bounds in this paper, and we have contented ourselves with the bounds of Theorem A which, on the other hand, are quite reasonable.

Theorem A follows without much effort from the following result, which yields structural information about the verbal subgroup $\omega(G)$ provided that it is soluble (equivalently, that G is soluble), without assuming that G_ω is finite.

Theorem B. *Let ω be an outer commutator word, and let G be a soluble group. Then there exists a series of subgroups from 1 to $\omega(G)$ such that:*

- (i) *All subgroups of the series are normal in G .*
- (ii) *Every section of the series is abelian and can be generated by values of ω all of whose powers are also values of ω in that section.*

Furthermore, the length of this series only depends on the word ω and on the derived length of G .

The existence of such a series was proved in [1] for derived words, and this particular case is the starting point for our proof of Theorem B. However, dealing with an arbitrary outer commutator word ω is a much more delicate matter, since one has to keep control of the nesting of commutators in ω , and then there might be problems such as the commutator of two values of ω not being necessarily a value of ω (contrary to the case of derived words). Our approach to the general case is geometric: we associate a labelled binary tree to every outer commutator word, a tree which reflects clearly the structure of the word, and which makes it easy to compare any two outer commutator words. Then the argument proceeds by measuring, with the help of the tree, how distant the word in question is from being a derived word, and using induction on this distance. The tree of an outer commutator word is introduced in Section 2, together with some related concepts that will be needed, and the proofs of Theorems A and B are postponed to Section 3.

We would like to remark that our proof of Theorem A is independent of and provides an alternative to the proof of the conciseness of outer commutator words given by Wilson in [9]. Wilson's argument is rather intricate and difficult to follow, and our geometric method provides a proof which, we honestly believe, is much easier to understand. Also, Wilson's proof goes by way of contradiction, and consequently he does not obtain any explicit bounds. On the other hand, notice that our proof lies within ZF, contrary to the proof of the existence of bounds for concise words via ultraproducts. To end this introduction, let us say that we are highly convinced that both the 'tree method' introduced in this paper and Theorem B may prove important tools for addressing other problems related to outer commutator words.

2. THE TREE OF AN OUTER COMMUTATOR WORD

As already mentioned, a fundamental device for the proof of Theorem B is to associate a labelled binary tree to every outer commutator word. For this purpose, we give a recursive and more formal definition of outer commutator words, and we use the same recursion to introduce the height and the labelled tree of such a word. In the following, we say that two words α and β are *disjoint* if the sets of indeterminates appearing in the two words are disjoint.

Definition 2.1. The set of outer commutator words, and the height and the labelled tree of an outer commutator word, are defined recursively as follows:

- (i) An indeterminate is an outer commutator of height 0, and its tree is an isolated vertex, labelled with the name of the indeterminate.
- (ii) If α and β are disjoint outer commutator words, then also $\omega = [\alpha, \beta]$ is an outer commutator word. The height $\text{ht}(\omega)$ of the word ω is taken to be the maximum of the heights of α and β plus 1, and the tree of ω is obtained by adding a new vertex with label ω and connecting it to the vertices labelled α and β of the corresponding trees of these words.

The tree of an outer commutator word ω provides a visual way of reading how ω is constructed by nesting commutators, easier than writing the actual expression of ω by using commutator brackets. We draw these trees by going downwards whenever we form a new commutator, so that the vertex with label ω is placed at the root of the tree. Every vertex v is labelled with an outer commutator word, which we denote by ω_v . Note that the indeterminates correspond exactly to the vertices of degree 1. Also, the height of ω coincides with the height of the tree, that is, the largest distance from the root to another vertex of the tree (which will be necessarily labelled by an indeterminate). For example, the following are the trees for the words γ_4 and δ_3 :




FIGURE 1. The trees of the words γ_4 and δ_3 .

More generally, the full tree of height h corresponds to the derived word δ_h .

All labels of the tree of an outer commutator word are completely determined, up to equivalence, by the tree itself (as a graph without labels): given the tree, we only need to associate an indeterminate to every vertex of degree 1, and then proceed downwards by labeling each vertex with the commutator of the labels of its immediate ascendants.

Observe that, if $\omega = [\alpha, \beta]$ is an outer commutator word, then the verbal subgroup $\omega(G)$ coincides with the commutator subgroup $[\alpha(G), \beta(G)]$.

If ω is an outer commutator word, then the set G_ω is clearly invariant under conjugation by elements of G . We remark that G_ω is not a subgroup in general; however, it has the following property.

Lemma 2.2. *Let ω be an outer commutator word. Then G_ω is symmetric, that is, $x \in G_\omega$ implies that $x^{-1} \in G_\omega$.*

Proof. We use induction on the height of ω . If $\omega = x_1$ then the result is true. Now assume that $\omega = [\alpha, \beta]$, where α, β are outer commutator words whose height is smaller than $\text{ht}(\omega)$. An element of G_ω is of the form $[y, z]$, with $y \in G_\alpha, z \in G_\beta$. Then $[y, z]^{-1} = [y^z, z^{-1}]$, where $y^z \in G_\alpha$ because G_α is invariant under conjugation, and $z^{-1} \in G_\beta$ by induction. So $[y, z]^{-1} \in G_\omega$, as we wanted to prove. $\square$

In the context of outer commutator words, in order to simplify the writing of words, it is convenient to reinterpret expressions such as $[\alpha, \alpha]$ (which is 1 in the free group to which α belongs), by replacing the second α by an equivalent word whose set of indeterminates is disjoint from that of the first α . More generally, we apply the same idea to every commutator $[\alpha, \beta]$ in which α and β have some indeterminate in common, so that $[\alpha, \beta]$ is a well-defined outer commutator word up to equivalence. Allowing this notation, the derived words can be defined by $\delta_0 = x_1$ and $\delta_i = [\delta_{i-1}, \delta_{i-1}]$ for $i \geq 1$, and the lower central words by $\gamma_1 = x_1$ and $\gamma_i = [\gamma_{i-1}, \gamma_1]$ for $i \geq 2$. Also, the tree corresponding to the word $[\gamma_3, \gamma_3]$ is the following:




FIGURE 2. The tree of the outer commutator word $[\gamma_3, \gamma_3]$.

We note that the vertices of the tree are naturally positioned in levels. More formally, we have the following.

Definition 2.3. Let v be a vertex of the tree of an outer commutator word ω of height h . We say that v is in the i -th level of the tree if it lies at distance $h - i$ from the root of the tree.

Thus the upmost level will be level 0 and the root will be at level h , but note that a vertex v at level i is not necessarily labelled with a word ω_v of height i , it might even happen that ω_v is an indeterminate.

It is also useful to associate a *companion* vertex to each vertex of the tree different from the root, defined as follows.

Definition 2.4. Let p be a vertex of the tree of an outer commutator word ω , different from the root, and let u be the immediate descendant of p . Then the *companion* of p is the only other vertex q of the tree which has u as an immediate descendant.

It is clear that companion vertices lie on the same level of the tree.

As said in the introduction, we will prove Theorem B for a general outer commutator word ω by induction on the ‘distance’ of ω to the closest derived word. We make this notion of distance precise in the following definition.

Definition 2.5. Let ω be an outer commutator word of height h . Then the *defect* of ω , which is denoted by $\text{def } \omega$, is defined as

$$\text{def } \omega = 2^{h+1} - 1 - V,$$

where V is the number of vertices of the tree of ω .

So, if the height of ω is h , then the defect is the number of vertices that need to be added to the tree of ω in order to get the tree of δ_h . Thus the defect is 0 if and only if ω is a derived word, and we have $\text{def } \gamma_4 = 8$ and $\text{def}[\gamma_3, \gamma_3] = 4$.

Let now φ and ω be two words, and let F be the free group to which φ belongs. We say that φ is ω -valued if $\varphi \in F_\omega$. If this is the case, then we have $G_\varphi \subseteq G_\omega$ for every group G , and in particular $\varphi(G) \leq \omega(G)$. For example, δ_2 is γ_3 -valued, but not conversely.

Definition 2.6. Let φ and ω be two outer commutator words. Then:

- (i) We say that ω is a *constituent* of φ if ω is, up to equivalence, the label of a vertex in the tree of φ .
- (ii) We say that φ is an *extension* of ω , or that ω is a *restriction* of φ , if the tree of φ is an upward extension of the tree of ω (simply as a tree, without labels).

Thus, in order to get an extension of ω , we only need to draw new binary trees at some of the vertices which are labelled by indeterminates in the tree of ω . Equivalently, a restriction of ω is obtained by selecting a number of vertices and erasing all branches lying on top of these vertices in the tree of ω .




FIGURE 3. An extension of $[\gamma_4, \delta_2]$.

In Figure 3, the black tree represents the word $\omega = [\gamma_4, \delta_2]$, and the extension of ω which is obtained by adding the grey trees is $\varphi = [[\gamma_3, \gamma_3], [\delta_2, \gamma_3]]$. Without having to check the commutator structure of these two words, the trees show that φ is ω -valued. On the other hand, observe that the derived word δ_h is an extension of all words of height less than or equal to h .

The following lemma is straightforward.

Lemma 2.7. Let φ and ω be two outer commutator words. Then:

- (i) If ω is a constituent of φ , then $\varphi(G) \leq \omega(G)$.
- (ii) If φ is an extension of ω , then φ is ω -valued.

3. PROOF OF THEOREMS A AND B

Before proceeding to the proof of Theorem B, we need some lemmas. First, we need to introduce the following concept.

Definition 3.1. Let T be the tree associated to an outer commutator word ω . A subset S of vertices of T is called a *section* of T if S is maximal (with respect to inclusion) subject to the condition that S does not contain two vertices which are one a descendant of the other. Equivalently, in terms of labels, this means that every indeterminate involved in ω appears in exactly one word ω_v with $v \in S$.

Visually, taking a section is nothing but cutting the tree from side to side.




FIGURE 4. A section of $[\gamma_3, \gamma_3, \delta_2]$.

A very natural way of obtaining a section is by cutting a tree below level i , that is, we consider the section S containing all vertices at level $i + 1$ and all the vertices of the tree lying below level $i + 1$ labelled by an indeterminate. This is the type of section that we will use in the proof of Theorem B.




FIGURE 5. Section of $[\gamma_4, \gamma_4]$ by cutting below level 0.

If $\omega = [\alpha, \beta]$ and γ are two outer commutator words, then by the Three Subgroup Lemma, we have

$$[\omega(G), \gamma(G)] \leq \pi^{(1)}(G)\pi^{(2)}(G),$$

where $\pi^{(1)} = [[\alpha, \gamma], \beta]$ and $\pi^{(2)} = [\alpha, [\beta, \gamma]]$ are also outer commutator words. Observe that the tree of $\pi^{(1)}$ is very similar to that of ω : one only needs to replace the tree on top of the vertex labelled α with the tree corresponding to $[\alpha, \gamma]$. The same happens with $\pi^{(2)}$, with β playing the role of α . The following lemma is a generalization of this fact; instead of stopping at the vertices labelled α and β , by iterating the process we can reach an arbitrary section of the tree.

Lemma 3.2. *Let ω be an outer commutator word, and let T be the tree of ω . If γ is another outer commutator word, then for every $v \in T$, we define $\pi^{(v)}$ to be the word whose tree is obtained by replacing the tree of ω_v at vertex v with the tree of $[\omega_v, \gamma]$. Then, for every section S of T , and for every group G , we have*

$$[\omega(G), \gamma(G)] \leq \prod_{v \in S} \pi^{(v)}(G).$$

Proof. We argue by induction on the number n of vertices of S . The case $n = 1$ is obvious, so we assume that $n \geq 2$. We also observe that the product $\prod_{v \in S} \pi^{(v)}(G)$ depends only on the subgroups $\pi^{(v)}(G)$, for $v \in S$, and not on the order in which they appear, since all those subgroups are normal in G . Let p be a vertex in S which has maximum distance from the root, let q be its companion vertex, and let u be the immediate descendant of p and q . Since each of the indeterminates involved in the word ω_u appears in exactly one of the words ω_v with $v \in S$, it necessarily follows from the assumption about p that $q \in S$. Now let S' be the section of T which is obtained from S by deleting p and q , and inserting u . By applying the induction hypothesis to S' , we have

$$(1) \quad [\omega(G), \gamma(G)] \leq \pi^{(u)}(G) \prod_{\substack{v \in S \\ v \neq p, q}} \pi^{(v)}(G).$$

On the other hand, by the Three Subgroup Lemma,

$$\begin{aligned} [\omega_u(G), \gamma(G)] &= [\omega_p(G), \omega_q(G), \gamma(G)] \\ &\leq [[\omega_p(G), \gamma(G)], \omega_q(G)][\omega_p(G), [\omega_q(G), \gamma(G)]], \end{aligned}$$

and consequently $\pi^{(u)}(G) \leq \pi^{(p)}(G)\pi^{(q)}(G)$, which completes the proof by (1). $\square$

Definition 3.3. Let ω be an outer commutator word, and let G be a group. A series of normal subgroups of G is said to be *power-closed generated* (or a *PCG-series*, for short) with respect to ω if every section H/J of the series is abelian and can be generated by values of ω in G/J all of whose powers are again values of ω in G/J .

It is clear that a series $H_0 \leq H_1 \leq \dots \leq H_n$ of normal subgroups of a group G is a PCG-series with respect to ω if and only if, for every $i = 1, \dots, n$, the quotient H_i/H_{i-1} is abelian and we can choose a subset S_i of G_ω such that:

- (P1) $H_i/H_{i-1} = \langle xH_{i-1} \mid x \in S_i \rangle$, i.e. $H_i = \langle S_i \rangle H_{i-1}$.
- (P2) $x^n H_{i-1} \in (G/H_{i-1})_\omega$ for every $x \in S_i$ and every $n \in \mathbb{Z}$.

Furthermore, since the set of ω -values is closed under conjugation, and the subgroups in a PCG-series are normal, we may assume if necessary that S_i is a normal subset of G .

Obviously, any PCG-series with respect to ω beginning from the trivial subgroup is contained in $\omega(G)$, and the content of Theorem B is precisely that, starting from 1, we can always reach $\omega(G)$ with a PCG-series provided that G is soluble.

Moreover, we note that if φ is another outer commutator word which is ω -valued, then any PCG-series with respect to φ is also a PCG-series with respect to ω . We will repeatedly use this fact in the sequel without further mention.

Now we state two more lemmas that we need for the proof of Theorem B.

Lemma 3.4. *Let ω be an outer commutator word, let G be a group, and let K and L be two normal subgroups of G . If there are two PCG-series with respect to ω from 1 to K and from 1 to L , then there is also a PCG-series from 1 to KL .*

Proof. If H/J is a normal abelian section of G generated by values of ω such that all of their powers are also values of ω , then also the section HL/JL has this property. Now the result readily follows. $\square$

Lemma 3.5. *Let α and β be two outer commutator words, and let G be a group. If there is a PCG-series from K to L in the group G with respect to α , and if $[L, \beta(G), L] = 1$, then by taking commutators with $\beta(G)$ we obtain a PCG-series from $[K, \beta(G)]$ to $[L, \beta(G)]$ with respect to $[\alpha, \beta]$. In particular, if there is a PCG-series from 1 to $\alpha(G)$ with respect to α , and if $[\alpha(G), \beta(G), \alpha(G)] = 1$, then there is also a series from 1 to $[\alpha(G), \beta(G)]$ with respect to $[\alpha, \beta]$.*

Proof. We first note that the condition $[L, \beta(G), L] = 1$ implies in particular that $[L, \beta(G)]$ is abelian, so that any section of this group is also abelian.

Let $K = H_0 \leq H_1 \leq \dots \leq H_n = L$ be a PCG-series with respect to α . We fix an integer i from 1 to n , and choose a normal subset S_i of G which is contained in G_α , and which satisfies properties (P1) and (P2) above. We claim that the set $T_i = \{[x, y] \mid x \in S_i, y \in G_\beta\}$ satisfies (P1) and (P2) for the section $[H_i, \beta(G)]/[H_{i-1}, \beta(G)]$ and the word $[\alpha, \beta]$. This proves the result, since T_i is contained in $G_{[\alpha, \beta]}$.

First of all, since S_i and G_β are normal subsets of G , the same is true about T_i . Then $N = \langle T_i \rangle [H_{i-1}, \beta(G)]$ is a normal subgroup of G , and H_i and $\beta(G)$ clearly commute modulo N . Thus $[H_i, \beta(G)] \leq N$, and property (P1) follows. Now let $[x, y]$ be an element of T_i , with $x \in S_i$ and $y \in G_\beta$. By using the fact that $[L, \beta(G), L] = 1$, we have $[x, y]^n = [x^n, y]$ for every $n \in \mathbb{Z}$. Since S_i satisfies (P2), we can write $x^n = a_n b_n$, with $a_n \in H_{i-1}$ and $b_n \in G_\alpha$. Thus

$$[x, y]^n = [a_n b_n, y] = [a_n, y]^{b_n} [b_n, y] \equiv [b_n, y] \pmod{[H_{i-1}, \beta(G)]},$$

which proves that (P2) holds for T_i . $\square$

Now we can easily see that Theorem B is true for derived words. This fact is already proved in Lemma 3.3 of [1], and the proof we provide is essentially the same. We include it here for the sake of completeness, and because the use of Lemma 3.5 simplifies the presentation. In the following, $G^{(i)}$ will denote as usual the i -th term $\delta_i(G)$ of the derived series of a group G .

Theorem 3.6. *Let G be a soluble group. Then, for every $i \geq 0$, there exists a PCG-series from 1 to $G^{(i)}$ with respect to δ_i . Furthermore, if the derived length of G is d , there is such a series of length at most $2^d - 2^i$ if $d \geq i$, or 0 if $d \leq i$.*

Proof. We first deal with the particular case when $G^{(i)}$ is abelian. Let us prove, by induction on i , that there is a PCG-series of length 2^i from 1 to $G^{(i)}$ with respect to δ_i . This is obvious for $i = 0$, so we assume that $i \geq 1$ and that the result holds for $i - 1$. If we apply it to the group G' , we obtain a PCG-series of length at most 2^{i-1} from 1 to $G^{(i)}$ with respect to δ_{i-1} . By Lemma 3.5, with $\alpha = \beta = \delta_{i-1}$, it follows that there is a PCG-series of the same length from 1 to $[G^{(i)}, G^{(i-1)}]$ with respect to δ_i . On the other hand, if we use again the result for δ_{i-1} , but in this case with the group $G/G^{(i)}$, we get a PCG-series of length at most 2^{i-1} from $G^{(i)}$ to $G^{(i-1)}$ with respect to δ_{i-1} . Another application of Lemma 3.5 yields a PCG-series from $[G^{(i)}, G^{(i-1)}]$ to $[G^{(i-1)}, G^{(i-1)}] = G^{(i)}$ with respect to δ_i . Now we can connect the two PCG-series with respect to δ_i that we have obtained so far, and the induction is complete.

Let us now deal with the general case. If $i \geq d$ there is nothing to prove, so we assume that $i < d$. By the last paragraph, for every j between i and $d - 1$ there is a PCG-series from $G^{(j+1)}$ to $G^{(j)}$ with respect to δ_j , of length 2^j . Since δ_j is δ_i -valued for $j \geq i$, by connecting these series we obtain a PCG-series from 1 to $G^{(i)}$ with respect to δ_i of length at most $2^d - 2^i$, as desired. $\square$

We can now prove Theorem B for arbitrary outer commutator words.

Proof of Theorem B. We concentrate on proving the existence of a PCG-series with respect to ω from 1 to $\omega(G)$; a close examination of the proof that follows shows that the length of the PCG-series constructed only depends on ω and on the derived length of G , and not on the particular group G .

We argue by double induction: we first use induction on the height of the word ω , and then, for a fixed value of the height, induction on the defect of ω . If ω has height 0, then $\omega = x_1$ and the result is trivially true. Now assume that $h = \text{ht}(\omega) \geq 1$ and that the result has been proved for any outer commutator word whose height is less than h . If $\text{def}(\omega) = 0$ then ω is a derived word, and the result holds by Theorem 3.6, so we assume that $\text{def}(\omega) > 0$. Let us write $\omega = [\alpha, \beta]$, where α and β are outer commutator words of height smaller than h . Then we have a PCG-series from 1 to $\alpha(G)$ with respect to α , and another one from 1 to $\beta(G)$ with respect to β . If we can reduce ourselves to the case that $[\omega(G), \alpha(G)] = 1$ or that $[\omega(G), \beta(G)] = 1$, then the proof of the theorem will be complete by invoking Lemma 3.5.

Let Φ be the (finite) set of all outer commutator words of height h which are a proper extension of ω . By the induction hypothesis on the defect, for every φ in Φ , there is a PCG-series from 1 to $\varphi(G)$ with respect to ω , since φ is ω -valued according to Lemma 2.7. By using Lemma 3.4, we can combine the series corresponding to all different words in Φ , and get a single PCG-series whose last term L contains $\varphi(G)$ for all φ in Φ . For the theorem to be proved, it suffices to find the desired PCG-series with respect to ω in the quotient G/L , and so we may assume in the remainder that $\varphi(G) = 1$ for all φ in Φ . We cannot guarantee in general that $[\omega, \alpha]$ or $[\omega, \beta]$ belong to the set Φ . However, we prove below that at least one of the subgroups

$[\omega(G), \alpha(G)]$ and $[\omega(G), \beta(G)]$ is contained in a product of verbal subgroups corresponding to words in Φ , and is consequently equal to 1, as desired.

Let i be the largest integer for which there is a vertex in the tree of ω at level i with label δ_i . Note that $1 \leq i < h$, since ω is not a derived word. Let S be the section of the tree of ω obtained by cutting the tree below level i , so that S contains all vertices at level $i + 1$ and all the vertices of the tree lying below level $i + 1$ which are labelled with an indeterminate. For every vertex v in S , we construct a word $\omega^{(v)}$ as follows. If the label ω_v of v is not an indeterminate, then we can write $\omega_v = [\omega_p, \omega_q]$, where p and q are the companion vertices at level i having v as immediate descendant. By the maximality of i , one of these vertices is labelled with a word which is different from δ_i . For simplicity, let us assume that this happens for q , the vertex on the right (the argument is exactly the same otherwise). We define $\omega^{(v)}$ to be the word whose tree is obtained by replacing ω_q with δ_i in the tree of ω . Thus the label of $\omega^{(v)}$ at the vertex v is the commutator $[\omega_p, \delta_i]$. On the other hand, if ω_v is an indeterminate, then $\omega^{(v)}$ is defined simply by putting the tree corresponding to δ_i on top of the vertex v in the tree of ω .




FIGURE 6. The two different cases for the construction of $\omega^{(v)}$ with $v \in S$. Observe that $i = 1$ in this example.

In any case, it is clear that $\text{ht}(\omega^{(v)}) = h$ and that $\omega^{(v)}$ is a proper extension of ω , so that $\omega^{(v)}$ belongs to Φ . Consequently, we have $\omega^{(v)}(G) = 1$ for every vertex v in the section S .

On the other hand, if we apply Lemma 3.2 to the section S with δ_i playing the role of γ , then we have

$$(2) \quad [\omega(G), \delta_i(G)] \leq \prod_{v \in S} \pi^{(v)}(G).$$

Here, $\pi^{(v)}$ is the word whose tree is obtained by inserting the tree of $[\omega_v, \delta_i]$ at vertex v in the tree of ω . Now, it is easy to compare the two words $\omega^{(v)}$ and $\pi^{(v)}$: they look the same at all vertices of the original tree of ω , except for the vertex v , where $\pi^{(v)}$ has the label $[\omega_v, \delta_i]$ and $\omega^{(v)}$ has either $[\omega_p, \delta_i]$ or δ_i . In any of the two cases, we have

$$(\pi^{(v)})_v(G) \leq (\omega^{(v)})_v(G),$$

and then, since $\pi^{(v)}$ and $\omega^{(v)}$ have the same labels outside the tree above v , also

$$\pi^{(v)}(G) \leq \omega^{(v)}(G).$$

Since this happens for all vertices in S , it follows from (2) that $[\omega(G), \delta_i(G)] = 1$. Now, by the definition of i , the derived word δ_i is a constituent of either

α or β , and consequently either $[\omega(G), \alpha(G)] = 1$ or $[\omega(G), \beta(G)] = 1$. As explained above, this completes the proof. $\square$

Finally, we derive Theorem A from Theorem B by adapting the argument given by Brazil, Krasilnikov and Shumyatsky in [1] for the case of derived words. We will need Dietzmann's Lemma, whose proof can be found in [5, 14.5.7].

Lemma 3.7. *If G is a group and $X = \{c_1, \dots, c_n\}$ is a normal subset, then every element $y \in \langle X \rangle$ is of the form $y = \prod_{i=1}^n c_i^{r_i}$, for some integers $r_1, \dots, r_n$.*

Proof of Theorem A. Suppose first that G is soluble. If $\omega(G) = 1$ the result is trivial, so we may assume that $\omega(G)$ is not the identity subgroup. By Theorem B, there is a PCG-series

$$1 = H_0 < H_1 < \dots < H_n = \omega(G).$$

Since G_ω is finite, each of the abelian quotients H_i/H_{i-1} can be generated by a finite number of values of ω all of whose powers are again values of ω . Then we can refine this PCG-series to a subnormal series

$$1 = G_0 < G_1 < \dots < G_k = \omega(G)$$

in which every section G_i/G_{i-1} is a non-trivial cyclic group consisting entirely of values of ω . Observe that, contrary to the original PCG-series, the length of this refined series may depend on the group G (more precisely, on the rank of G); however, this will have no effect in the proof. Now, for every non-trivial element x in G_i/G_{i-1} , there exists $y \in G_\omega \setminus \{1\}$ such that $x = yG_{i-1}$, and consequently

$$|G_1/G_0| + |G_2/G_1| + \dots + |G_k/G_{k-1}| \leq m + k - 1.$$

Observe that $\log_2 |G_i/G_{i-1}| \leq |G_i/G_{i-1}| - 1$ for all i , since $|G_i/G_{i-1}| \geq 2$. Hence

$$|\omega(G)| = \prod_{i=1}^k |G_i/G_{i-1}| = 2^{\sum_{i=1}^k \log_2 |G_i/G_{i-1}|} \leq 2^{\sum_{i=1}^k |G_i/G_{i-1}| - k} \leq 2^{m-1},$$

which proves part (i) of Theorem A.

Now assume that G is non-soluble. Observe that $m \geq 3$ in this case, since otherwise $\omega(G)$ is cyclic and G is soluble. If h is the height of ω , then δ_h is ω -valued, and the same holds for δ_{h+1} . Let $|G_{\delta_{h+1}}| = l$. Then $|(G/G^{(h+1)})_\omega| \leq m - l + 1$, and by the bound for the soluble case, it follows that

$$|\omega(G)/G^{(h+1)}| \leq 2^{m-l}.$$

Now we bound the order of $G^{(h+1)}$. We claim that the order of an element $g \in G_{\delta_{h+1}}$ is at most $(m-1)(m-2)$. Of course, we may assume $g \neq 1$. Let us write $g = [a, b]$ with $a, b \in G_{\delta_h}$, and consider the subgroup $H = \langle a, b \rangle$. Let $C = C_H(a)$. Since $a \in G_\omega \setminus \{1\}$, it has at most $m-1$ conjugates in G , and consequently $|H : C| \leq m-1$. Now C permutes the $m-1$ non-trivial values of ω in G , and leaves the element a fixed by definition. Thus $|C : C_C(b)| \leq m-2$, and consequently $|H : Z(H)| = |H : C_H(a) \cap C_H(b)| \leq$

$(m-1)(m-2)$. By applying Schur's Theorem [5, 10.1.4] to H , it follows that the exponent of H' is at most $(m-1)(m-2)$, which proves the claim.

Let $G_{\delta_{h+1}} = \{c_0, c_1, c_2, \dots, c_{l-1}\}$, where $c_0 = 1$. By Lemma 2.2, the set $G_{\delta_{h+1}}$ is symmetric, so we can assume that $c_i = c_i^{-1}$ for every $i = 0, \dots, t$ for some $0 \leq t \leq l-1$ and that $c_{t+2j} = c_{t+2j-1}^{-1}$ for each $j = 1, \dots, (l-1-t)/2$ (note that $l-1-t$ is even). Since $G_{\delta_{h+1}}$ is a normal subset of G , it follows from Lemma 3.7 that every element w of $\delta_{h+1}(G)$ is of the form $c_1^{n_1} c_2^{n_2} \cdots c_{l-1}^{n_{l-1}}$, where $1 \leq n_i \leq |c_i|$. Now, we have two choices for each n_i with $1 \leq i \leq t$ (if $t \geq 1$, otherwise we have nothing to choose) and at most $(m-1)(m-2)$ choices for each product of the form $c_{t+2j-1}^{n_{t+2j-1}} c_{t+2j}^{n_{t+2j}} = c_{t+2j-1}^{n_{t+2j-1}-n_{t+2j}}$. So

$$|G^{(h+1)}| \leq 2^t [(m-1)(m-2)]^{\frac{l-1-t}{2}} \leq 2^t (m-1)^{l-1-t} \leq (m-1)^{l-1},$$

and we conclude that

$$\begin{aligned} |\omega(G)| &= |\omega(G)/G^{(h+1)}| |G^{(h+1)}| \leq 2^{m-l} (m-1)^{l-1} \\ &\leq (m-1)^{m-1}, \end{aligned}$$

since $m \geq 3$. This completes the proof of Theorem A. $\square$

4. APPENDIX: EXISTENCE OF BOUNDS VIA ULTRAPRODUCTS

In this appendix, we give two different proofs of the following result, mentioned in the introduction.

Theorem 4.1. *Let ω be a concise word. Then, there exists a function $f : \mathbb{N} \rightarrow \mathbb{N}$ such that, if G is a group in which $|G_\omega| \leq m$, then $|\omega(G)| \leq f(m)$.*

For the convenience of the reader, we begin by recalling briefly the construction of ultraproducts of groups. To this end, we need the concept of an ultrafilter. (See [2] for an account on ultraproducts from an algebraic point of view.)

Definition 4.2. A *filter* over a non-empty set I is a non-empty family $\mathcal{U}$ of subsets of I such that:

- (i) The intersection of two elements of $\mathcal{U}$ also lies in $\mathcal{U}$.
- (ii) If P is in $\mathcal{U}$ and $P \subseteq Q$, then also Q is in $\mathcal{U}$.
- (iii) The empty set does not belong to $\mathcal{U}$.

The filter $\mathcal{U}$ is called *principal* if it consists of all supersets of a fixed subset of I , and it is called an *ultrafilter* if it is maximal in the set of all filters over I ordered by inclusion.

Equivalently, a filter $\mathcal{U}$ over I is an ultrafilter if and only if, for every subset J of I , either $J \in \mathcal{U}$ or $I \setminus J \in \mathcal{U}$. By (i) and (iii) above, only one of these conditions holds.

The existence of non-principal ultrafilters is independent of the Zermelo-Fraenkel axioms for set theory. It can be easily proved by using the Axiom of Choice, but is in fact weaker than that. On the other hand, an ultrafilter over I is non-principal if and only if it contains all cofinite subsets of I .

Definition 4.3. Let I be a non-empty set, and let $\mathcal{U}$ be an ultrafilter over I . The *ultraproduct* modulo $\mathcal{U}$ of a family $\mathcal{G} = \{G_i\}_{i \in I}$ of groups is the quotient of the cartesian product $\prod_{i \in I} G_i$ (i.e. the unrestricted direct product) by the subgroup consisting of all tuples $(g_i)_{i \in I}$ such that the set

$$\{i \in I \mid g_i = 1\}$$

lies in $\mathcal{U}$. We denote this ultraproduct by $\mathcal{G}_{\mathcal{U}}$.

Thus two tuples $(g_i)_{i \in I}$ and $(h_i)_{i \in I}$ of the cartesian product define the same element of the ultraproduct $\mathcal{G}_{\mathcal{U}}$ if and only if the set of indices i for which $g_i = h_i$ lies in $\mathcal{U}$. In the remainder of the paper, we use the bar notation for the image of an element or a subset of $\prod_{i \in I} G_i$ in an ultraproduct.

The first proof of Theorem 4.1 that we present is based on the following particular case of Łoś's Theorem from model theory. (See Theorem 3.1 and Corollary 3.2 of [2].)

Lemma 4.4. *Let $\mathcal{G} = \{G_i\}_{i \in I}$ be a family of groups and let $\mathcal{U}$ be an ultrafilter over I . Then, a sentence in the first-order language of groups holds in the ultraproduct $\mathcal{G}_{\mathcal{U}}$ if and only if the set of all $i \in I$ for which the sentence holds in G_i is a member of $\mathcal{U}$.*

Recall that the *width* of a word ω in a group G is the supremum, as g ranges over the verbal subgroup $\omega(G)$, of the minimum length of all decompositions of g as a product of elements of $G_{\omega} \cup G_{\omega}^{-1}$. Obviously, if G is finite, then ω has finite width in G . We may similarly speak of the width of ω over a subset S of $\omega(G)$, by taking the supremum only over elements of S .

First proof of Theorem 4.1. This proof is based on the following two facts:

(i) For a given positive integer m , the property that ω takes at most m values in a group can be expressed as a sentence in the first-order language of groups. More precisely, if ω involves k indeterminates, we may use the following formula:

$$\exists g_{11} \dots \exists g_{1k} \dots \exists g_{m1} \dots \exists g_{mk} \forall x_1 \dots \forall x_k \bigvee_{i=1}^m \omega(x_1, \dots, x_k) = \omega(g_{i1}, \dots, g_{ik}).$$

(ii) For a given positive integer n , the property that ω has width at most n in a group can be expressed as a sentence in the first-order language of groups. To see this, note that this property is equivalent to every product of $n+1$ elements of $G_{\omega} \cup G_{\omega}^{-1}$ being also a product of n elements of that set.

Assume, by way of contradiction, that there is an infinite sequence G_n of groups such that $|(G_n)_{\omega}| \leq m$ for all $n \in \mathbb{N}$ but $|\omega(G_n)|$ goes to infinity. Choose a non-principal ultrafilter $\mathcal{U}$, and let $Q = \mathcal{G}_{\mathcal{U}}$. Then, by Lemma 4.4 and (i), we have $|Q_{\omega}| \leq m$. It follows that $|\omega(Q)|$ is finite, since ω is concise. Then ω has finite width, say k , in Q . By Lemma 4.4 again, this time used together with (ii), there is a subset $J \in \mathcal{U}$ such that ω has width at most k in G_n for all $n \in J$. Since $|(G_n)_{\omega}| \leq m$, it follows that $|\omega(G_n)| \leq (2m)^k$ for every $n \in J$. This is incompatible with the condition $\lim_{n \rightarrow \infty} |\omega(G_n)| = \infty$: since $\mathcal{U}$ is a non-principal ultrafilter, every cofinite subset of $\mathbb{N}$ has non-empty intersection with J . $\square$

Now we give a second proof of Theorem 4.1, which only needs the definition and basic properties of ultraproducts, and which is independent of Łoś's Theorem. This proof basically follows an argument communicated to us by Avinoam Mann.

Lemma 4.5. *Let $\mathcal{G} = \{G_n\}_{n \in \mathbb{N}}$ be a family of groups, and for every $n \in \mathbb{N}$, let S_n be a non-empty finite subset of G_n . If $\mathcal{U}$ is an ultrafilter over $\mathbb{N}$ then the cardinality of the image of $S = \prod_{n \in \mathbb{N}} S_n$ in the ultraproduct $\mathcal{G}_{\mathcal{U}}$ is given by*

$$(3) \quad |\overline{S}| = \sup_{J \in \mathcal{U}} \left(\min_{n \in J} |S_n| \right),$$

provided that the supremum is finite, and $\overline{S}$ is infinite otherwise. In particular:

- (i) *If $|S_n| \leq k$ for all n , then $|\overline{S}| \leq k$.*
- (ii) *If the ultrafilter $\mathcal{U}$ is non-principal and $|S_n| \geq k$ for big enough n , then $|\overline{S}| \geq k$.*

Proof. Let J be an arbitrary element of $\mathcal{U}$, and put $m = \min_{n \in J} |S_n|$. Let us prove that $|\overline{S}| \geq m$, which gives one of the inequalities in (3). For every $n \in \mathbb{N}$, we consider m elements $s_n^{(1)}, \dots, s_n^{(m)} \in S_n$, which we take different if $n \in J$ and arbitrary if $n \notin J$. Let

$$s^{(i)} = (s_n^{(i)})_{n \in \mathbb{N}}, \quad \text{for every } i = 1, \dots, m.$$

We claim that the images of $s^{(i)}$ and $s^{(j)}$ in $\mathcal{G}_{\mathcal{U}}$ are different for all $i \neq j$. Otherwise, the tuples $s^{(i)}$ and $s^{(j)}$ coincide on a subset $X \in \mathcal{U}$, but they are different by construction on $J \in \mathcal{U}$. Hence $J \subseteq \mathbb{N} \setminus X$ and, by (ii) of the definition of a filter, we also have $\mathbb{N} \setminus X \in \mathcal{U}$. Thus both X and $\mathbb{N} \setminus X$ lie in $\mathcal{U}$, which is impossible since $\mathcal{U}$ is an ultrafilter. This proves our claim, and consequently that $|\overline{S}| \geq m$. Observe that this also proves that $\overline{S}$ is infinite if the supremum in (3) is not finite.

For the reverse inequality, put $r = \sup_{J \in \mathcal{U}} (\min_{n \in J} |S_n|)$, and assume that r is finite. By way of contradiction, suppose that $|\overline{S}| \geq r + 1$. If $s^{(1)}, \dots, s^{(r+1)}$ are elements of S whose images in $\mathcal{G}_{\mathcal{U}}$ are all different, then for all $i, j \in \{1, \dots, r+1\}$, $i \neq j$, the set

$$X_{ij} = \{n \in \mathbb{N} \mid s_n^{(i)} \neq s_n^{(j)}\}$$

belongs to $\mathcal{U}$. Hence the intersection J of all the X_{ij} is also in $\mathcal{U}$. Now observe that, if $n \in J$, then $s_n^{(1)}, \dots, s_n^{(r+1)}$ are all different and, consequently, $|S_n| \geq r + 1$. It follows that $\min_{n \in J} |S_n| \geq r + 1$, which is a contradiction with the definition of r .

Finally, observe that (i) is obvious, and that (ii) follows because a non-principal ultrafilter contains all cofinite subsets. $\square$

If ω is a word and $\{G_i\}_{i \in I}$ is an infinite family of groups, it is not always the case that $\omega(\prod_{i \in I} G_i) = \prod_{i \in I} \omega(G_i)$, and only the inclusion $\subseteq$ may be guaranteed. Our next lemma is an approximation to the reverse inclusion.

Lemma 4.6. *Let ω be a word, and let $\{G_i\}_{i \in I}$ be a family of groups. Suppose that $S_i \subseteq \omega(G_i)$ for every $i \in I$, and that the width of ω can be uniformly*

bounded over all the subsets S_i . Then

$$\prod_{i \in I} S_i \subseteq \omega\left(\prod_{i \in I} G_i\right).$$

Proof. Let $g = (g_i)_{i \in I} \in \prod_{i \in I} S_i$. If the width of ω is at most k over all the subsets S_i , then every g_i can be written as a product of k elements $x_i^{(1)}, \dots, x_i^{(k)}$ of $(G_i)_\omega \cup (G_i)_\omega^{-1}$. We use these elements to define $2k$ elements of G_i as follows: for every $j = 1, \dots, k$, we put

$$g_i^{(2j-1)} = \begin{cases} x_i^{(j)}, & \text{if } x_i^{(j)} \in (G_i)_\omega, \\ 1, & \text{otherwise,} \end{cases} \quad \text{and} \quad g_i^{(2j)} = \begin{cases} 1, & \text{if } x_i^{(j)} \in (G_i)_\omega, \\ x_i^{(j)}, & \text{otherwise.} \end{cases}$$

Then $g_i^{(2j-1)} \in (G_i)_\omega$, $g_i^{(2j)} \in (G_i)_\omega^{-1}$ and $g_i = g_i^{(1)} \dots g_i^{(2k)}$ for every $i \in I$. If we put $g^{(r)} = (g_i^{(r)})_{i \in I}$ for $r = 1, \dots, 2k$, it follows that $g^{(2j-1)} \in (\prod_{i \in I} G_i)_\omega$ and $g^{(2j)} \in (\prod_{i \in I} G_i)_\omega^{-1}$ for $j = 1, \dots, k$, and also that $g = g^{(1)} \dots g^{(2k)}$. Thus $g \in \omega(\prod_{i \in I} G_i)$, as desired. $\square$

Lemma 4.7. *Let ω be a word, and let G be a group such that $|\omega(G)| \geq k$, where k is a positive integer. Then, there exists a subset S of $\omega(G)$ such that $|S| \geq k$ and ω has width less than k over S .*

Proof. For every integer $i \geq 0$, let T_i be the subset of all elements of $\omega(G)$ of (minimum) length i with respect to the set of generators $G_\omega \cup G_\omega^{-1}$. Put $T = \cup_{i=0}^{k-1} T_i$. If T_i is non-empty for every $i = 0, \dots, k-1$, then $|T| \geq k$ and we may take $S = T$. If, on the contrary, T_i is empty for some $i = 0, \dots, k-1$, then ω has width at most $i-1$ in G , and then we may take $S = \omega(G)$. $\square$

Second proof of Theorem 4.1. By way of contradiction, assume that there is a family $\{G_n\}_{n \in \mathbb{N}}$ of groups such that $|(G_n)_\omega| \leq m$ for all n , but nevertheless $\lim_{n \rightarrow \infty} |\omega(G_n)| = \infty$. Let us fix an arbitrary positive integer k . According to Lemma 4.7, if n is big enough, there is a subset S_n of $\omega(G_n)$ such that $|S_n| \geq k$ and ω has width less than k over S_n . We complete the sequence $\{S_n\}_{n \in \mathbb{N}}$ by choosing the first terms equal to 1. Now, if $G = \prod_{n \in \mathbb{N}} G_n$ and $S = \prod_{n \in \mathbb{N}} S_n$, we have

$$G_\omega = \prod_{n \in \mathbb{N}} (G_n)_\omega, \quad \text{and} \quad S \subseteq \omega(G),$$

where the last inclusion follows from Lemma 4.6. Consider now a non-principal ultrafilter $\mathcal{U}$ over $\mathbb{N}$, and let $Q = \mathcal{G}_{\mathcal{U}}$ be the corresponding ultraproduct. Then $Q_\omega = \overline{(G_\omega)}$ and $\omega(Q) = \overline{\omega(G)} \supseteq \overline{S}$. By applying Lemma 4.5, we obtain that $|Q_\omega| \leq m$ and $|\omega(Q)| \geq k$. Since k is arbitrary, we get $|\omega(Q)| = \infty$, which is a contradiction, since the word ω is concise. $\square$

Acknowledgments. We want to thank Avinoam Mann for communicating to us the argument we have used in the second proof of Theorem 4.1, and for giving us permission to include it in this paper. On the other hand, we want to express our gratitude to the University of the Basque Country and the University of Padova for their hospitality while this work was carried out.

REFERENCES

- [1] S. Brazil, A. Krasilnikov, P. Shumyatsky, *Groups with bounded verbal conjugacy classes*, J. Group Theory **9** (2006), 127–137.
- [2] P.C. Eklof, *Ultraproducts for algebraists*, Chapter A.3 in J. Barwise (editor), *Handbook of Mathematical Logic*, Studies in Logic and the Foundations of Mathematics **90**, North-Holland, Amsterdam, 1982, pp. 105–137.
- [3] S.V. Ivanov, *P. Hall's conjecture on the finiteness of verbal subgroups*, Izv. Vyssh. Ucheb. Zaved. **325** (1989), 60–70.
- [4] P.M. Neumann, M.R. Vaughan-Lee, *An essay on BFC groups*, Proc. London Math. Soc. (3) **35** (1977), 213–237.
- [5] D.J. Robinson, *A course in the theory of groups*, 2nd edition, Springer-Verlag, Berlin, 1996.
- [6] D. Segal, A. Shalev, *On groups with bounded conjugacy classes*, Quart. J. Math. Oxford (2) **50** (1999), 505–516.
- [7] R.F. Turner-Smith, *Marginal subgroup properties for outer commutator words*, Proc. London Math. Soc. **14** (1964), 321–341.
- [8] R.F. Turner-Smith, *Finiteness conditions for verbal subgroups*, J. London Math. Soc. **41** (1966), 166–176.
- [9] J. Wilson, *On outer-commutator words*, Canadian J. Math. **26** (1974), 608–620.

MATEMATIKA SAILA, EUSKAL HERRIKO UNIBERTSITATEA, 48080 BILBAO, SPAIN
E-mail address: `gustavo.fernandez@ehu.es`

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI BOLOGNA, PIAZZA DI PORTA SAN
DONATO 5, 40127 BOLOGNA, ITALY
E-mail address: `mmorigi@dm.unibo.it`